



# Altered metabolic requirements in cancer cell migration and metastasis

## Citation

Lee, Jaewon, M Rosa Ng, Kerstin W Sinkevicius, Carla F Kim, Gaudenz Danuser, Joan S Brugge, and Marcia C Haigis. 2012. Altered metabolic requirements in cancer cell migration and metastasis. BMC Proceedings 6(Suppl 3): P32.

## Published Version

doi:10.1186/1753-6561-6-S3-P32

## Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:10436271>

## Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

## Share Your Story

The Harvard community has made this article openly available.  
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

POSTER PRESENTATION

Open Access

# Altered metabolic requirements in cancer cell migration and metastasis

Jaewon Lee<sup>1,2\*</sup>, M Rosa Ng<sup>1</sup>, Kerstin W Sinkevicius<sup>3,4,5</sup>, Carla F Kim<sup>3,4,5</sup>, Gaudenz Danuser<sup>1</sup>, Joan S Brugge<sup>1</sup>, Marcia C Haigis<sup>1,2</sup>

From Metabolism, diet and disease  
Washington, DC, USA. 29-31 May 2012

## Background

Metastasis poses a great challenge in clinical management of many cancers. Metabolic demands of cancer cell proliferation – i.e. elevated aerobic glycolysis for biomass generation – have been well characterized, but the contribution of altered metabolism to metastasis remains to be elucidated. While elevated aerobic glycolysis, a phenomenon termed Warburg effect, is a hallmark of proliferative tumor cells, emerging evidence suggests that metastatic cancer cells have an opposite phenotype.

## Materials and methods

We investigated the metabolism in cell lines of different metastatic capacity. In addition, to quantitatively study cell migration and metabolism, we used MCF10A breast epithelial cells with fluorescent histone tags along with live cell imaging and tracking by a custom MATLAB program to measure speed and behavior of migration.

## Results

Our preliminary data suggest that the more metastatic cancer cells depend on mitochondrial metabolism. Furthermore, changing mitochondrial metabolism in MCF10A breast epithelial cell line affected not only the speed but also the pattern of cell migration. Consistent with the altered migratory behaviors, stimulation of mitochondrial metabolism changed cell adhesion markers.

## Conclusions

In sum, we show evidence that mitochondrial metabolism plays an important role in promoting cell migration

and altering cell adhesion with implications for cancer metastasis.

## Author details

<sup>1</sup>Department of Cell Biology, Harvard Medical School, Boston, MA 02115, USA. <sup>2</sup>The Paul F. Glenn Labs for the Biological Mechanisms of Aging, Harvard Medical School, Boston, MA 02115, USA. <sup>3</sup>Stem Cell Program, Children's Hospital Boston, Boston, MA 02115, USA. <sup>4</sup>Department of Genetics, Harvard Medical School, Boston, MA 02115, USA. <sup>5</sup>Harvard Stem Cell Institute, Cambridge, MA 02138, USA.

Published: 1 June 2012

doi:10.1186/1753-6561-6-S3-P32

**Cite this article as:** Lee et al.: Altered metabolic requirements in cancer cell migration and metastasis. *BMC Proceedings* 2012 6(Suppl 3):P32.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



<sup>1</sup>Department of Cell Biology, Harvard Medical School, Boston, MA 02115, USA

Full list of author information is available at the end of the article